

# Metric approach to quantum constraints

Dorje C Brody<sup>1</sup>, Anna C T Gustavsson<sup>2</sup>, and  
Lane P Hughston<sup>1</sup>

<sup>1</sup> Department of Mathematics, Imperial College London, London SW7 2AZ, UK

<sup>2</sup> Blackett Laboratory, Imperial College London, London SW7 2AZ, UK

**Abstract.** A new framework for deriving equations of motion for constrained quantum systems is introduced, and a procedure for its implementation is outlined. In special cases the framework reduces to a quantum analogue of the Dirac theory of constraints in classical mechanics. Explicit examples involving spin- $\frac{1}{2}$  particles are worked out in detail: in one example our approach coincides with a quantum version of the Dirac formalism, while the other example illustrates how a situation that cannot be treated by Dirac's approach can nevertheless be dealt with in the present scheme.

Submitted to: *J. Phys. A: Math. Gen.*

## 1. Introduction

Recently there has been a renewed interest in understanding the properties of constrained quantum dynamics [1, 2, 3, 4]. The key idea behind quantum constraints is the fact that the space of pure states (rays through the origin of Hilbert space) is a symplectic manifold, and hence that Dirac's theory of constraints [5, 6] in classical mechanics is applicable in the quantum regime. The quantum state space is also equipped with a metric structure—generally absent in a classical phase space—induced by the probabilistic features of quantum mechanics. In the context of analysing constrained quantum motions it is therefore natural to examine the theory from the viewpoint of metric geometry, as opposed to a treatment based entirely on symplectic geometry. This is the goal of the present paper.

The metric approach that we propose is not merely a reformulation of the Dirac formalism using the quantum symplectic structure. Indeed, there are two distinct advantages in the metric approach over the symplectic approach: (a) the metric approach to quantum constraints is in general more straightforward to implement, even in situations when the constraints can be treated by the symplectic method; and (b) there are nontrivial examples of constraints that cannot be implemented in the symplectic approach but can be implemented in the metric approach. Our plan therefore is first to outline the general metric approach to quantum constraints, and

then to consider specific examples. We also derive and examine a necessary condition for the metric approach to be equivalent to the symplectic formalism. The first example examined here concerns the system consisting of a pair of spin- $\frac{1}{2}$  particles. We impose the constraint that the state should lie on the product subspace of the total state space, upon which all the energy eigenstates lie. This is the example considered in [1, 4] using the symplectic approach. Here we analyse the problem using the metric approach, and show that the constrained equations of motion reduce to those obtained in [4]. The second example concerns a single spin- $\frac{1}{2}$  particle, and we impose the constraint that an observable that does not commute with the Hamiltonian must be conserved. This is perhaps the simplest example of a quantum constraint that is not evidently tractable in the symplectic approach but can be readily dealt with by use of the metric approach.

## 2. Geometry of quantum state space

We begin by remarking that the space of pure quantum states associated with a Hilbert space of dimension  $n$  is the projective Hilbert space  $\mathcal{P}^{n-1}$  of dimension  $n - 1$  (see [7, 8, 9, 10, 11] and references cited therein). We regard  $\mathcal{P}^{n-1}$  as a real even-dimensional manifold  $\Gamma$ , and denote a typical point in  $\Gamma$ , corresponding to a ray in the associated Hilbert space, by  $\{x^a\}_{a=1,2,\dots,2n-2}$ . It is well known that  $\Gamma$  has an integrable complex structure. Since the complex structure of  $\Gamma$  plays an important role in what follows, it may be helpful if we make a few general remarks about the relevant ideas.

We recall that an even-dimensional real manifold  $\mathfrak{M}$  is said to have an almost complex structure if there exists a global tensor field  $J_b^a$  satisfying

$$J_c^a J_b^c = -\delta_b^a, \quad (1)$$

The almost complex structure is then said to be integrable if the Nijenhuis tensor

$$N_{ab}^c = J_d^c \nabla_{[a} J_{b]}^d - J_{[a}^d \nabla_{|d|} J_{b]}^c \quad (2)$$

vanishes [12]. It is straightforward to check that  $N_{ab}^c$  is independent of the choice of symmetric connection  $\nabla_a$  on  $\mathfrak{M}$ . The vanishing of  $N_{ab}^c$  can be interpreted as follows. A complex vector field on  $\mathfrak{M}$  is said to be of positive (*resp.*, negative) type if  $J_b^a V^b = +iV^a$  (*resp.*,  $J_b^a V^b = -iV^a$ ). The vanishing of  $N_{ab}^c$  is a necessary and sufficient condition for the commutator of two vector fields of the same type to be of that type.

A Riemannian metric  $g_{ab}$  on  $\mathfrak{M}$  is said to be compatible with an almost complex structure  $J_b^a$  if the following conditions hold: (i) the metric is Hermitian:

$$J_c^a J_d^b g_{ab} = g_{cd}, \quad (3)$$

and (ii) the almost complex structure is covariantly constant:

$$\nabla_a J_c^b = 0, \quad (4)$$

where  $\nabla_a$  is the torsion-free Riemannian connection associated with  $g_{ab}$ . An alternative expression for the Hermitian condition is that  $\Omega_{ab} = -\Omega_{ba}$ , where

$$\Omega_{ab} = J_a^c g_{bc}. \quad (5)$$

It follows that  $\nabla_a J_c^b = 0$  if and only if  $\nabla_a \Omega_{bc} = 0$ . However, if the almost complex structure is integrable, then a sufficient condition for  $\nabla_a J_c^b = 0$  is  $\nabla_{[a} \Omega_{bc]} = 0$ . This follows on account of the identity

$$\nabla_a \Omega_{bc} = \frac{3}{2} J_b^p J_c^q \nabla_{[a} \Omega_{pq]} - \frac{3}{2} \nabla_{[a} \Omega_{bc]} + \frac{1}{4} \Omega_{ad} N_{bc}^d. \quad (6)$$

A manifold  $\mathfrak{M}$  with an integrable complex structure and a compatible Riemannian structure is called a Kähler manifold. The antisymmetric tensor  $\Omega_{ab}$  is then referred to as the ‘fundamental two-form’ or Kähler two-form. It follows from the definition of  $\Omega_{ab}$  along with the Hermitian condition on  $g_{ab}$  that  $\Omega_{ab}$  itself is Hermitian in the sense that

$$J_c^a J_d^b \Omega_{ab} = \Omega_{cd}. \quad (7)$$

Furthermore, we find the tensor  $\Omega^{ab}$  defined by using the inverse metric to raise the indices of  $\Omega_{ab}$ , so

$$\Omega^{ab} = g^{ac} g^{db} \Omega_{cd}, \quad (8)$$

acts as an inverse to  $\Omega_{ab}$ . In particular, we have

$$\Omega^{ac} \Omega_{bc} = \delta_b^a. \quad (9)$$

In the case of quantum theory there is a natural Riemannian structure on the manifold  $\Gamma$ , called the Fubini-Study metric. If  $x$  and  $y$  represent a pair of points in  $\Gamma$ , and  $|\psi(x)\rangle$  and  $|\psi(y)\rangle$  are representative Hilbert space vectors, then the Fubini-Study distance between  $x$  and  $y$  is given by  $\theta$ , where

$$\frac{\langle \psi(y) | \psi(x) \rangle \langle \psi(x) | \psi(y) \rangle}{\langle \psi(x) | \psi(x) \rangle \langle \psi(y) | \psi(y) \rangle} = \frac{1}{2} (1 + \cos \theta). \quad (10)$$

The Kähler form  $\Omega_{ab}$  can be used to define a one-parameter family of symplectic structures on  $\Gamma$ , given by  $\kappa \Omega_{ab}$ , where  $\kappa$  is a nonvanishing real constant. In quantum mechanics the symplectic structure defined by

$$\omega_{ab} = \frac{1}{2} \Omega_{ab} \quad (11)$$

plays a special role. In particular, if we define the inverse symplectic structure by  $\omega^{ab} = 2\Omega^{ab}$  so that  $\omega^{ac} \omega_{bc} = \delta_b^a$ , and if we choose units such that  $\hbar = 1$ , then we find that the Schrödinger trajectories on  $\Gamma$  are given by Hamiltonian vector fields of the form

$$\dot{x}^a = \omega^{ab} \nabla_b H, \quad (12)$$

where

$$H(x) = \frac{\langle \psi(x) | \hat{H} | \psi(x) \rangle}{\langle \psi(x) | \psi(x) \rangle}. \quad (13)$$

Thus we see that the expectation of the Hamiltonian operator  $\hat{H}$  gives rise to a real function  $H(x)$  on  $\Gamma$ . This function plays the role of the Hamiltonian in the determination of the symplectic flow associated with the Schrödinger trajectory.

### 3. Metric formalism for quantum constraints

With these geometric tools in hand we now proceed to formulate the metric approach to quantum constraints. We assume that the system under investigation is subject to a family of  $N$  constraints

$$\Phi^i(x) = 0, \quad (14)$$

where  $i = 1, \dots, N$ . The condition  $\Phi^i(x) = 0$  for each  $i$  defines a hypersurface in  $\Gamma$ . The intersection of all  $N$  such hypersurfaces then defines the constraint subspace in  $\Gamma$  onto which the dynamics must be restricted. To enforce the constraint in the metric approach we remove from the vector field  $\dot{x}^a$  those components that are normal to the constraint subspace. The equations of motion then take the form

$$\dot{x}^a = \omega^{ab} \nabla_b H - \lambda_i g^{ab} \nabla_b \Phi^i, \quad (15)$$

where  $\omega^{ab}$  is the inverse quantum symplectic structure, and where  $\{\lambda_i\}_{i=1,\dots,N}$  are the Lagrange multipliers associated with the constraints. In order to determine  $\{\lambda_i\}$  we consider the identity  $\dot{\Phi}^j = 0$ . Using the chain rule this can be expressed as

$$\dot{x}^a \nabla_a \Phi^j = 0. \quad (16)$$

Substitution of (15) in (16) then gives

$$\omega^{ab} \nabla_a \Phi^j \nabla_b H - \lambda_i g^{ab} \nabla_a \Phi^j \nabla_b \Phi^i = 0. \quad (17)$$

To simplify (17) we define the symmetric matrix

$$M^{ij} = g^{ab} \nabla_a \Phi^i \nabla_b \Phi^j. \quad (18)$$

If the matrix  $M^{ij}$  is nonsingular, we write  $M_{ij}$  for its inverse such that  $M_{ik} M^{kj} = \delta_i^j$ . We can then solve (17) for  $\lambda_i$  to find

$$\lambda_i = M_{ij} \omega^{ab} \nabla_a \Phi^j \nabla_b H. \quad (19)$$

Substituting the expression for  $\lambda_i$  back into (15) we find that the constrained equations of motion are given by

$$\dot{x}^a = \omega^{ab} \nabla_b H - M_{ij} \omega^{cd} \nabla_c \Phi^j \nabla_d H g^{ab} \nabla_b \Phi^i. \quad (20)$$

This is the main result of the paper.

We remark that in the case for which the system of constraints  $\{\Phi^i\}$  corresponds to a family of quantum observables,  $M^{ij}$  for fixed  $i, j$  represents the covariance between the two observables  $\hat{\Phi}^i$  and  $\hat{\Phi}^j$  in the state represented by the point  $x$ :

$$M^{ij} = \langle \hat{\Phi}^i \hat{\Phi}^j \rangle - \langle \hat{\Phi}^i \rangle \langle \hat{\Phi}^j \rangle. \quad (21)$$

It follows that if there is a single constraint for a conserved observable (as in Example 2 below), then  $M$  is never singular except at isolated points of zero measure because the variance of an observable is positive except at its eigenstates. In the case of two conserved observables, say,  $\hat{\Phi}^1 = \hat{A}$  and  $\hat{\Phi}^2 = \hat{B}$ , we have

$$M^{ij} = \begin{pmatrix} \text{var}(\hat{A}) & \text{cov}(\hat{A}, \hat{B}) \\ \text{cov}(\hat{A}, \hat{B}) & \text{var}(\hat{B}) \end{pmatrix}. \quad (22)$$

The determinant  $\Delta$  of  $M^{ij}$  is then given by

$$\Delta = (1 - \rho^2) \text{var}(\hat{A}) \text{var}(\hat{B}), \quad (23)$$

where  $\rho$  is the correlation of  $\hat{A}$  and  $\hat{B}$ . Since the variances are positive (except at eigenstates),  $\Delta$  can identically vanish if and only if  $\rho = \pm 1$ , i.e. either  $\hat{A}$  and  $\hat{B}$  are perfectly correlated or they are perfectly anticorrelated. Analogous observations can be made when there are more than two conserved observables for the constraint, and we conclude that  $M^{ij}$  is singular only when the constraints are redundant. Therefore, in the case of constraints on conserved observables the metric approach introduced here is always implementable, provided that we do not introduce redundant constraints.

In the case of algebraic constraints for which the constraint system  $\{\Phi^i\}$  does not correspond to a family of quantum observables (as in Example 1 below), we are unaware of any physical characteristic that might lead to a singular behaviour in  $M^{ij}$ . Thus in this case the invertibility of  $M^{ij}$  must be examined individually.

#### 4. Equivalence of metric and symplectic approaches

Before considering specific examples of constrained systems that can be described using the present approach, it will be of interest to ask how this framework might be related to the approach of Dirac [5, 6], or more precisely, its quantum counterpart [1, 2, 4] which we shall refer to as the symplectic approach. In the symplectic approach, the constrained equations of motion can be expressed in the same form as (12), but with a modified inverse symplectic structure  $\tilde{\omega}^{ab}$  in place of  $\omega^{ab}$ , which in effect is the induced symplectic form on the constraint surface [4]. Thus, we would like to know under what condition the metric approach leading to the right side of (20) also reduces to a modified symplectic flow of the form  $\tilde{\omega}^{ab} \nabla_b H$ , where  $H$  is the same Hamiltonian as the one in the original Schrödinger equation (12). Intuitively, we would expect that when there is an even number of constraints, the two methods might become equivalent. However, the verification or falsification of this assertion turns out to be difficult. Nevertheless, in what follows we shall establish the sufficient condition under which the symplectic approach and the metric approach become equivalent.

In order to investigate whether (20) can be rewritten in the form

$$\dot{x}^a = \tilde{\omega}^{ab} \nabla_b H \quad (24)$$

for a suitably defined antisymmetric tensor  $\tilde{\omega}^{ab}$ , we rearrange terms in (20) to write

$$\begin{aligned} \dot{x}^a &= (\omega^{ad} - M_{ij} g^{ab} \nabla_b \Phi^i \omega^{cd} \nabla_c \Phi^j) \nabla_d H \\ &= (\omega^{ad} - \mu_{bc} g^{ab} \omega^{cd}) \nabla_d H, \end{aligned} \quad (25)$$

where we have defined

$$\mu_{bc} = M_{ij} \nabla_b \Phi^i \nabla_c \Phi^j. \quad (26)$$

It follows that we need to find whether the expression

$$\tilde{\omega}^{ab} = \omega^{ab} - g^{ad} \omega^{cb} \mu_{dc} \quad (27)$$

defines a symplectic structure on the subspace of the state space. Since the symplectic structure  $\omega^{ab}$  is antisymmetric, we shall examine under which condition  $\tilde{\omega}^{ab}$  is antisymmetric. This is equivalent to asking whether the following relation holds:

$$g^{ac}\omega^{bd}\mu_{cd} \stackrel{?}{=} -g^{bc}\omega^{ad}\mu_{cd}. \quad (28)$$

Suppose that (28) is valid. Then using (5) and (11) we can rewrite (28) as

$$g^{ac}g^{be}J_e^d\mu_{cd} = -g^{bc}g^{ae}J_e^d\mu_{cd}. \quad (29)$$

Transvecting this with  $g_{fa}$  and relabelling the indices we obtain

$$g^{be}J_e^d\mu_{ad} = -g^{bc}J_a^d\mu_{cd}, \quad (30)$$

from which it follows that

$$J_b^d\mu_{ad} = -J_a^d\mu_{bd}. \quad (31)$$

Multiplying both sides of (31) with  $J_c^a$  we find that the condition (28) is equivalent to

$$J_a^cJ_b^d\mu_{cd} = \mu_{ab}, \quad (32)$$

where we have used (1) and the symmetry of  $\mu_{ab}$ . It follows that for the metric formulation of the constraint motion (20) to be expressible in the Hamiltonian form (24) with the original Hamiltonian  $H$  the matrix  $\mu_{bc}$  defined by (26) must be Hermitian. This is the sufficiency condition that we set out to establish. Let us now examine specific cases to gain insights into this condition.

*Single constraint case.* In general the  $J$ -invariance condition (32) need not be satisfied. One can easily see this by considering the case for which there is only one constraint given by  $\Phi(x) = 0$ . Then expression (18) becomes a scalar quantity, which we will denote by  $M$ , and thus its inverse is  $M^{-1}$ . It follows that

$$\mu_{ab} = M^{-1}\nabla_a\Phi\nabla_b\Phi. \quad (33)$$

Substituting this expression for  $\mu_{ab}$  into (32) gives us

$$J_a^c\nabla_c\Phi J_b^d\nabla_d\Phi = \nabla_a\Phi\nabla_b\Phi, \quad (34)$$

which implies that

$$J_a^c\nabla_c\Phi = \nabla_a\Phi, \quad (35)$$

but this clearly is a contradiction, since the two vectors  $J_a^c\nabla_c\Phi$  and  $\nabla_a\Phi$  are *orthogonal*. We see therefore that in the case of a single constraint the condition (32) is not satisfied, and the constraint motion (20) cannot be expressed in the Hamiltonian form (24).

*Two constraint case.* Let us examine the case in which there are two constraints. As we shall show, in this case the  $J$ -invariance condition for  $\mu_{bc}$  reduces to a simpler condition. Let us write  $\Phi^1 = A$  and  $\Phi^2 = B$  for the two constraints. Then the inverse of the matrix  $M^{ij}$  can be written in the form

$$M_{ij} = \frac{1}{2\Delta} \epsilon_{ii'}\epsilon_{jj'}M^{i'j'}, \quad (36)$$

where  $\Delta = \det(M^{ij})$ , and  $\epsilon_{ij}$  is a totally-skew tensor with  $i, i', j, j' = 1, 2$ . Substituting (36) into (26) we find

$$\begin{aligned}\mu_{bc} &= M_{ij} \nabla_b \Phi^i \nabla_c \Phi^j \\ &= \frac{1}{2\Delta} \epsilon_{ii'} \epsilon_{jj'} g^{pq} \nabla_p \Phi^{i'} \nabla_q \Phi^{j'} \nabla_b \Phi^i \nabla_c \Phi^j \\ &= \frac{1}{2\Delta} \epsilon_{ii'} \nabla_b \Phi^i \nabla_p \Phi^{i'} \epsilon_{jj'} \nabla_c \Phi^j \nabla_q \Phi^{j'} g^{pq} \\ &= \frac{1}{2\Delta} \tau_{bp} \tau_{cq} g^{pq},\end{aligned}\tag{37}$$

where we have defined

$$\tau_{ab} = \epsilon_{ij} \nabla_a \Phi^i \nabla_b \Phi^j.\tag{38}$$

In order for (37) to satisfy (32) we thus require that

$$J_a^b \tau_{bc} = \pm \tau_{ab} J_c^b,\tag{39}$$

where we have substituted (37) into condition (32) and used  $J_p^b J_q^c g^{pq} = g^{bc}$ . Since

$$\tau_{ab} = \nabla_a A \nabla_b B - \nabla_a B \nabla_b A,\tag{40}$$

we find that (39) can be written more explicitly in the form

$$J_a^b \nabla_b A \nabla_c B - J_a^b \nabla_b B \nabla_c A = \pm (\nabla_a A J_c^b \nabla_b B - \nabla_a B J_c^b \nabla_b A).\tag{41}$$

Hence if this condition is satisfied (with either a plus or a minus sign) by the two constraints  $A$  and  $B$ , then the constrained equations of motion (20) take the form (24).

*Holomorphic constraints.* If we choose the constraint function to be holomorphic so that the two constraints  $A$  and  $B$  are given by the real and imaginary parts of

$$\Phi(x) = A(x) + iB(x),\tag{42}$$

then we can find the form of  $\tau_{ab}$  for which (39) holds. To obtain an expression for  $\tau_{ab}$ , we recall that a real vector on  $\Gamma$  can be decomposed into its complex ‘positive’ and ‘negative’ parts  $V_a = (V_\alpha, V_{\alpha'})$  (cf. [13]). These components are given respectively by

$$(V_\alpha, 0) = \frac{1}{2}(V_a - iJ_a^b V_b), \quad \text{and} \quad (0, V_{\alpha'}) = \frac{1}{2}(V_a + iJ_a^b V_b).\tag{43}$$

Hence,  $V_{\alpha'}$  is the complex conjugate of  $V_\alpha$ , and these components are the eigenvectors of the complex structure  $J_b^a$  with eigenvalues  $\pm i$ . It follows that if the vector is of type  $V_a = (V_\alpha, 0)$  then  $J_b^a V_a = iV_b$ , and similarly if  $V_a = (0, V_{\alpha'})$  then  $J_b^a V_a = -iV_b$ . With respect to this decomposition the tensor  $\tau_{ab}$  can be expressed as

$$\tau_{ab} = \begin{pmatrix} \tau_{\alpha\beta} & \tau_{\alpha\beta'} \\ \tau_{\alpha'\beta} & \tau_{\alpha'\beta'} \end{pmatrix}.\tag{44}$$

We rewrite the condition (39) in the form

$$J_a^c J_b^d \tau_{cd} = \pm \tau_{ab}\tag{45}$$

by contracting both sides with  $J_d^c$  and using (1). In terms of the decomposition (44) we find that in order for (45) to be true we require that  $\tau_{ab}$  takes either of the two forms:

$$\tau_{ab}^{(+)} = \begin{pmatrix} 0 & \tau_{\alpha\beta'} \\ \tau_{\alpha'\beta} & 0 \end{pmatrix}, \quad \text{or} \quad \tau_{ab}^{(-)} = \begin{pmatrix} \tau_{\alpha\beta} & 0 \\ 0 & \tau_{\alpha'\beta'} \end{pmatrix}, \quad (46)$$

where the plus and minus in  $\tau_{ab}^{(\pm)}$  correspond to the required sign in (45). In view of (42) we can write the two constraints as  $A(x) = \frac{1}{2}(\Phi + \bar{\Phi})$  and  $B(x) = -\frac{1}{2}i(\Phi - \bar{\Phi})$ , where  $\bar{\Phi}$  denotes the complex conjugate of  $\Phi$ . Then we have:

$$\begin{aligned} \nabla_\alpha A &= \frac{1}{2}\nabla_\alpha \Phi, & \nabla_\alpha B &= -\frac{1}{2}i\nabla_\alpha \Phi, \\ \nabla_{\alpha'} A &= \frac{1}{2}\nabla_{\alpha'} \bar{\Phi}, & \nabla_{\alpha'} B &= \frac{1}{2}i\nabla_{\alpha'} \bar{\Phi}. \end{aligned} \quad (47)$$

Using these expressions together with (40) we find that the components of  $\tau_{ab}^{(-)}$  all vanish. Hence, when the two constraints  $A$  and  $B$  are given by (42), the condition (39) is satisfied when  $\tau_{ab}$  takes the form

$$\tau_{ab} = \begin{pmatrix} 0 & \frac{1}{2}i\nabla_\alpha \Phi \nabla_{\beta'} \bar{\Phi} \\ -\frac{1}{2}i\nabla_{\alpha'} \bar{\Phi} \nabla_\beta \Phi & 0 \end{pmatrix}, \quad (48)$$

i.e. when  $\tau_{ab} = \tau_{ab}^{(+)}$ . It follows that the metric formalism associated with holomorphic constraints of the form (42) is equivalent to the symplectic formalism.

We remark that the quadratic form  $\tilde{\omega}^{ab}$  acting from the right annihilates the vector  $\nabla_a \Phi^k$  normal to the constraint surface. This can be verified explicitly as follows:

$$\begin{aligned} \tilde{\omega}^{ad} \nabla_a \Phi^k &= \omega^{ad} \nabla_a \Phi^k - M_{ij} g^{ab} \nabla_b \Phi^i \nabla_a \Phi^k \omega^{cd} \nabla_c \Phi^j \\ &= \omega^{ad} \nabla_a \Phi^k - M_{ij} M^{ki} \omega^{cd} \nabla_c \Phi^j \\ &= \omega^{ad} \nabla_a \Phi^k - \omega^{ad} \nabla_a \Phi^k \\ &= 0, \end{aligned} \quad (49)$$

since  $g^{ab} \nabla_a \Phi^k \nabla_b \Phi^i = M^{ki}$  and  $M_{ij} M^{ki} = \delta_j^k$ . This condition, of course, is equivalent to the condition that  $\dot{x}^a \nabla_a \Phi^k = 0$ . However, in general we have

$$\tilde{\omega}^{ad} \nabla_a \Phi^k \neq 0, \quad (50)$$

since  $\tilde{\omega}^{ab}$  is not in general antisymmetric. In other words, the vanishing of the left side of (50) is equivalent to the  $J$ -invariance condition for  $\mu_{bc}$ .

In summary, the procedure for deriving the equations of motion in the metric approach to constrained quantum motion is as follows. First, express the relevant constraints in the form  $\{\Phi^i(x) = 0\}_{i=1,2,\dots,N}$ . Determine the matrix  $M^{ij}$  via (18). Assuming that  $M^{ij}$  is nonsingular, calculate its inverse  $M_{ij}$ . Substitute the result in (20), and we recover the relevant equations of motion. Having obtained the general procedure, let us now examine some explicit examples implementing this procedure.

## 5. Illustrative examples

*Example 1.* The first example that we consider here is identical to the one considered in [4] involving a pair of spin- $\frac{1}{2}$  particles. We consider the subspace of the state space



associated with product states upon which all the energy eigenstates lie. An initial state that lies on this product space is required to remain a product state under the evolution generated by a generic Hamiltonian. In this example there are two constraints  $\Phi^1(x)$  and  $\Phi^2(x)$ , and we shall show that the metric approach introduced here gives rise to a result that agrees with the one obtained in [4] using the symplectic approach.

Let us work with the coordinates of the quantum state space given by the ‘action-angle’ variables [4, 14], where the canonical conjugate variables are given by  $\{x^a\} = \{q_\nu, p_\nu\}_{\nu=1,\dots,n-1}$  such that when a generic pure states  $|x\rangle$  is expanded in terms of the energy eigenstates  $\{|E_\alpha\rangle\}_{\alpha=1,\dots,n}$ , the associated amplitudes are given by  $\{p_\nu\}$ , and the relative phases by  $\{q_\nu\}$ . In the case of a pair of spin- $\frac{1}{2}$  particles we can thus expand a generic state in the form

$$|x\rangle = \sqrt{p_1}e^{-iq_1}|E_1\rangle + \sqrt{p_2}e^{-iq_2}|E_2\rangle + \sqrt{p_3}e^{-iq_3}|E_3\rangle + \sqrt{1-p_1-p_2-p_3}|E_4\rangle. \quad (51)$$

This choice of coordinates has the property that if we write

$$H(p, q) = \frac{\langle x|H|x\rangle}{\langle x|x\rangle} \quad (52)$$

for the Hamiltonian, where  $|x\rangle = |p, q\rangle$ , then the Schrödinger equation is expressed in the form of the conventional Hamilton equations:

$$\dot{q}_\nu = \frac{\partial H(q, p)}{\partial p_\nu} \quad \text{and} \quad \dot{p}_\nu = -\frac{\partial H(q, p)}{\partial q_\nu}. \quad (53)$$

In the energy basis the Hamiltonian can be expressed as

$$H = \sum_{\alpha=1}^4 E_\alpha |E_\alpha\rangle\langle E_\alpha|. \quad (54)$$

It follows that the phase space function  $H(p, q)$  for a generic Hamiltonian is given by

$$H = E_4 + \sum_{\nu=1}^3 \Omega_\nu p_\nu, \quad (55)$$

where  $\Omega_\nu = E_\nu - E_4$ . The symplectic structure and its inverse in these coordinate are thus given respectively by

$$\omega_{ab} = \begin{pmatrix} \mathbb{O} & \mathbb{1} \\ -\mathbb{1} & \mathbb{O} \end{pmatrix} \quad \text{and} \quad \omega^{ab} = \begin{pmatrix} \mathbb{O} & \mathbb{1} \\ -\mathbb{1} & \mathbb{O} \end{pmatrix}, \quad (56)$$

where  $\mathbb{1}$  is the  $2 \times 2$  identity matrix and  $\mathbb{O}$  is the  $2 \times 2$  null matrix.

Let us write  $\{\psi^\alpha\}_{\alpha=1,2,3,4}$  for the coefficients of the energy eigenstates in (51). Then the constraint equation is expressed in the form  $\psi^1\psi^4 = \psi^2\psi^3$ , which is just a single complex equation [4]. Thus in real terms we have two constraints given by

$$\begin{cases} \Phi^1 = \sqrt{p_1 p_4} \cos q_1 - \sqrt{p_2 p_3} \cos(q_2 + q_3) = 0, \\ \Phi^2 = \sqrt{p_1 p_4} \sin q_1 - \sqrt{p_2 p_3} \sin(q_2 + q_3) = 0. \end{cases} \quad (57)$$

These constraints are ‘separable’ and can be rewritten as

$$\begin{cases} \Phi^1 = q_1 - q_2 - q_3 = 0, \\ \Phi^2 = p_1(1 - p_1 - p_2 - p_3) - p_2 p_3 = 0. \end{cases} \quad (58)$$

The next step in the metric approach for the constraint is to work out the expression for the Fubini-Study metric associated with the line element:

$$ds^2 = 8 \left[ \frac{\psi^{[\alpha} d\psi^{\beta]} \bar{\psi}_{[\alpha} d\bar{\psi}_{\beta]}}{(\bar{\psi}_\gamma \psi^\gamma)^2} \right] \quad (59)$$

in terms of the canonical variables  $\{q_\nu, p_\nu\}_{\nu=1,2,3}$ . A calculation shows that

$$g_{ab} = \begin{pmatrix} 4(1-p_1)p_1 & -4p_1p_2 & -4p_1p_3 & 0 & 0 & 0 \\ -4p_1p_2 & 4(1-p_2)p_2 & -4p_2p_3 & 0 & 0 & 0 \\ -4p_1p_3 & -4p_2p_3 & 4(1-p_3)p_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1-p_2-p_3}{p_1p_4} & \frac{1}{p_4} & \frac{1}{p_4} \\ 0 & 0 & 0 & \frac{1}{p_4} & \frac{1-p_1-p_3}{p_2p_4} & \frac{1}{p_4} \\ 0 & 0 & 0 & \frac{1}{p_4} & \frac{1}{p_4} & \frac{1-p_1-p_2}{p_3p_4} \end{pmatrix}, \quad (60)$$

where for simplicity we have denoted  $p_4 = 1 - p_1 - p_2 - p_3$ . The inverse is thus:

$$g^{ab} = \begin{pmatrix} \frac{1-p_2-p_3}{4p_1p_4} & \frac{1}{4p_4} & \frac{1}{4p_4} & 0 & 0 & 0 \\ \frac{1}{4p_4} & \frac{1-p_1-p_3}{4p_2p_4} & \frac{1}{4p_4} & 0 & 0 & 0 \\ \frac{1}{4p_4} & \frac{1}{4p_4} & \frac{1-p_1-p_2}{4p_3p_4} & 0 & 0 & 0 \\ 0 & 0 & 0 & (1-p_1)p_1 & -p_1p_2 & -p_1p_3 \\ 0 & 0 & 0 & -p_1p_2 & (1-p_2)p_2 & -p_2p_3 \\ 0 & 0 & 0 & -p_1p_3 & -p_2p_3 & (1-p_3)p_3 \end{pmatrix}. \quad (61)$$

To obtain explicit expressions for equations of motion (20) we need to calculate the matrix  $M^{ij}$  and its inverse. A short calculation shows that

$$M^{ij} = \begin{pmatrix} \frac{1}{4} \left( \frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} + \frac{1}{p_4} \right) & 0 \\ 0 & p_1p_4(1-4p_1p_4+4p_2p_3) \\ & + (p_2+p_3-4p_2p_3)(p_2p_3-p_1p_4) \end{pmatrix}, \quad (62)$$

from which its inverse  $M_{ij}$  can easily be obtained. Putting these together, we find that the equations of motion are given by

$$\begin{aligned} \dot{q}_1 &= \Omega_1 - \frac{p_2p_3(1-2p_1-p_2-p_3)(\Omega_1-\Omega_2-\Omega_3)}{p_2p_3(1-p_2-p_3)-p_1^2(p_2+p_3)+p_1(1-p_2-p_3)(p_2+p_3)} \\ \dot{q}_2 &= \Omega_2 + \frac{p_1p_3(1-p_1-p_3)(\Omega_1-\Omega_2-\Omega_3)}{p_2p_3(1-p_2-p_3)-p_1^2(p_2+p_3)+p_1(1-p_2-p_3)(p_2+p_3)} \\ \dot{q}_3 &= \Omega_3 + \frac{p_1p_2(1-p_1-p_2)(\Omega_1-\Omega_2-\Omega_3)}{p_2p_3(1-p_2-p_3)-p_1^2(p_2+p_3)+p_1(1-p_2-p_3)(p_2+p_3)} \\ \dot{p}_1 &= 0 \\ \dot{p}_2 &= 0 \\ \dot{p}_3 &= 0. \end{aligned} \quad (63)$$

We can simplify the equations using the relation  $p_1p_4 = p_2p_3$ , which gives us

$$\begin{aligned} \dot{q}_1 &= \Omega_1 - (1-2p_1-p_2-p_3)(\Omega_1-\Omega_2-\Omega_3) \\ \dot{q}_2 &= \Omega_2 + (p_1+p_3)(\Omega_1-\Omega_2-\Omega_3) \\ \dot{q}_3 &= \Omega_3 + (p_1+p_2)(\Omega_1-\Omega_2-\Omega_3) \\ \dot{p}_1 &= 0 \\ \dot{p}_2 &= 0 \\ \dot{p}_3 &= 0. \end{aligned} \quad (64)$$

These equations are precisely the ones obtained in [4] by means of the symplectic formalism. The solution to these equations are also worked out in [4].

Before we proceed to the next example, let us verify explicitly that the condition (32) for the equivalence of the metric and symplectic approaches indeed holds in this example. For this we need to obtain the expression for the complex structure in the canonical coordinates  $\{q_\nu, p_\nu\}_{\nu=1,2,3}$  using the relations (5) and (11). This is given by

$$J_b^a = \begin{pmatrix} 0 & 0 & 0 & \frac{1-p_2-p_3}{2p_1p_4} & \frac{1}{2p_4} & \frac{1}{2p_4} \\ 0 & 0 & 0 & \frac{1}{2p_4} & \frac{1-p_1-p_3}{2p_2p_4} & \frac{1}{2p_4} \\ 0 & 0 & 0 & \frac{1}{2p_4} & \frac{1}{2p_4} & \frac{1-p_1-p_2}{2p_3p_4} \\ 2(p_1-1)p_1 & 2p_1p_2 & 2p_1p_3 & 0 & 0 & 0 \\ 2p_1p_2 & 2(p_2-1)p_2 & 2p_2p_3 & 0 & 0 & 0 \\ 2p_1p_3 & 2p_2p_3 & 2(p_3-1)p_3 & 0 & 0 & 0 \end{pmatrix}, \quad (65)$$

where as before we write  $p_4 = 1 - p_1 - p_2 - p_3$ . Substituting this expression and the expression for  $\mu_{ab}$  obtained from (26) into (32), we find that the condition is indeed satisfied.

*Example 2.* Consider a single spin- $\frac{1}{2}$  particle system immersed in a  $z$ -field with a unit strength. The space of pure states for this system is just the surface of the Bloch sphere. The Hamiltonian of the system is given by  $\hat{H} = \hat{\sigma}_z$ , where  $\hat{\sigma}_z$  is the Pauli spin matrix in the  $z$ -direction. We then impose the constraint that an observable, say  $\hat{\sigma}_x$ , that does not commute with the Hamiltonian, must be conserved under the time evolution.

As before we chose the canonical coordinates  $x^a = \{q, p\}$  for the Bloch sphere by setting

$$|x(p, q)\rangle = \sqrt{1-p}|E_1\rangle + \sqrt{p}e^{-iq}|E_2\rangle. \quad (66)$$

The Hamiltonian in this coordinate system is given by

$$H(q, p) = 1 - 2p. \quad (67)$$

The conservation of  $\hat{\sigma}_x$  then reduces to a single real constraint of the form  $\langle x|\hat{\sigma}_x|x\rangle = \text{constant}$ , which, by use of (66), gives us

$$\Phi(x) = 2\sqrt{p(1-p)}\cos q. \quad (68)$$

The metric on the Bloch sphere, in terms of our conjugate variables, is given by


$$g_{ab} = \begin{pmatrix} 4(1-p)p & 0 \\ 0 & \frac{1}{(1-p)p} \end{pmatrix}. \quad (69)$$

The symplectic structure has the same form as (56) so we can now calculate the scalar quantity  $M$  defined in (18). A short calculation shows that

$$M = (1-2p)^2\cos^2 q + \sin^2 q. \quad (70)$$

Substituting (70) into (20) we find that the equations of motion are given by

$$\dot{q} = \frac{-2(1-2p)^2\cos^2 q}{(1-2p)^2\cos^2 q + \sin^2 q}, \quad (71)$$



**Figure 1.** (colour online) A field plot of the dynamics resulting from a system constrained to remain on a surface defined by  $\Phi = \hat{\sigma}_x$  when the system evolves according to the Hamiltonian  $H = \hat{\sigma}_z$ , where  $\hat{\sigma}_x$  and  $\hat{\sigma}_z$  are Pauli matrices. Shown in yellow are the integral curves of the motion when we chose as the starting positions  $\{\theta_1, \phi_n\} = \{\frac{23\pi}{48}, \frac{2\pi n}{24}\}_{n=0, \dots, 23}$  and  $\{\theta_2, \phi_n\} = \{\frac{25\pi}{48}, \frac{2\pi n}{24}\}_{n=0, \dots, 23}$ . The great circles shown in red, the equators of the  $x$  and  $z$ -axis, consist of fixed points. A state that initially lies on one of these great circles does not move away from that point, whereas all other states evolve asymptotically towards the fixed point where the associated integral curve intersects the equator of the sphere.

and

$$\dot{p} = \frac{4(1-2p)(-1+p)p \sin q \cos q}{(1-2p)^2 \cos^2 q + \sin^2 q}. \quad (72)$$

In order to visualise the results we convert the equations of motion into angular coordinates, using the method outlined in [4]. In angular coordinates (66) is given by

$$|x(\theta, \phi)\rangle = \cos \frac{1}{2}\theta |E_1\rangle + \sin \frac{1}{2}\theta e^{i\phi} |E_2\rangle. \quad (73)$$

Comparing (66) and (73) we identify  $p = \sin^2 \frac{1}{2}\theta$  and  $q = -\phi$ . The equations of motion then become:

$$\dot{\theta} = \frac{1}{2} \left( \frac{\sin(2\theta) \sin(2\phi)}{1 - \sin^2 \theta \cos^2 \phi} \right), \quad \text{and} \quad \dot{\phi} = \frac{2 \cos^2 \theta \cos^2 \phi}{1 - \sin^2 \theta \cos^2 \phi}. \quad (74)$$

Figure 1 shows some of the integral curves resulting from the above equations, plotted on the surface of the Bloch sphere. Equation (74) is valid at all points except where

$(\theta, \phi) = (\frac{\pi}{2}, 0)$  and where  $(\theta, \phi) = (\frac{\pi}{2}, \pi)$ . At these singular fixed points the method used cannot be defined.

It is also straightforward to verify that the results above could not have been obtained using the symplectic approach. The complex structure on the underlying state space of the spin- $\frac{1}{2}$  particle in our coordinate system is given by

$$J_b^a = \begin{pmatrix} 0 & \frac{1}{2(1-p)p} \\ -2(1-p)p & 0 \end{pmatrix}. \quad (75)$$

Evaluating (26) using (70), and substituting the result along with (75) into (32), we find that (32) does not hold.

- [1] Burić, N. 2007 Hamiltonian quantum dynamics with separability constraints. *Ann. Phys.* **323** 17–33
- [2] Corichi, A. 2008 On the geometry of quantum constrained systems. *Class. Quantum Grav.* **25** 135013.
- [3] Graefe, E. M., Korsch, H. J. & Niederle, A. E. 2008 Mean-field dynamics of a non-Hermitian Nose-Hubbard dimer. *Phys. Rev. Lett.* **101**, 150408.
- [4] Brody, D. C., Gustavsson, A. C. T. & Hughston, L. P. 2008 Symplectic approach to quantum constraints. *J. Phys. A: Math. Theor.* **41** 475301
- [5] Dirac, P. A. M. 1950 Generalized Hamiltonian dynamics. *Canadian J. Math.* **2**, 129–148.
- [6] Dirac, P. A. M. 1958, Generalised Hamiltonian dynamics. *Proc. R. Soc. London A* **246**, 326–332.
- [7] Kibble, T. W. B. 1979 Geometrisation of quantum mechanics. *Commun. Math. Phys.* **65**, 189–201.
- [8] Cirelli, R, Mania, A, Pizzocchero, L. 1990 Quantum mechanics as an infinite-dimensional Hamiltonian system with uncertainty structure. I, II. *J. Math. Phys.* **31**, 2891–2903.
- [9] Hughston, L. P. 1995 Geometric aspects of quantum mechanics. In *Twistor Theory* ed S Huggett (New York: Marcel Dekker, Inc.).
- [10] Ashtekar, A. & Schilling, T. A. 1995 Geometry of quantum mechanics. CAM-94 Physics Meeting, in AIP Conf. Proc. **342**, 471–478, ed. Zapeda, A. (AIP Press, Woodbury, New York).
- [11] Brody, D. C. & Hughston, L. P. 2001 Geometric quantum mechanics. *J. Geom. Phys.* **38**, 19–53.
- [12] Newlander, A & Nirenberg, L. 1957 Complex analytic coordinates in almost complex manifolds *Ann. Math.* **65**, 391–404.
- [13] Brody, D. C. & Hughston, L. P. 1998 Statistical geometry in quantum mechanics. *Proc. R. Soc. London A* **454**, 2445–2475.
- [14] Oh, P. & Kim, M. 1994 Action angle variables for complex projective space and semiclassical exactness. *Mod. Phys. Lett. A* **9**, 3339–3346.